

August 25, 1936

Prepared for
Am. Fish. Soc.

REPORT NO. 359A

STUDIES ON MERGANSER DEPREDACTIONS IN MICHIGAN TROUT WATERS

J. W. Leonard and David S. Shetter

Institute for Fisheries Research

Michigan Department of Conservation

This paper is a report on findings obtained from an analysis of the stomach contents of ninety-eight American Mergansers and two Red-breasted Mergansers, collected in pursuance of the Merganser Control Program of the Michigan Department of Conservation.

It is generally agreed that mergansers seldom concentrate heavily on trout streams except when severe cold prevails, and ice formation drives them from their preferred feeding areas in the bays and estuaries of the Great Lakes and the larger inland lakes. Michigan's present merganser policy recommends that control measures be adopted when census work reveals a concentration of more than twenty-five mergansers per mile of stream. Such control is customarily exercised by Conservation Officers.

The authors devoted some time during January, February and March, 1936 to stream patrol in various sections of the northern part of the lower peninsula, with a view to determining centers of merganser concentration. These field observations, coupled with distributional data afforded by collections made by conservation officers, indicate that the large rivers support a much heavier concentration of mergansers than do the small, fast trout streams. An exception to this is offered by the Platte River, in Benzie County, where the largest single collection of recovered ducks was made. This case may well be explained by the fact that the collection was made at a point within fifteen miles of Lake Michigan. Most of the trout water patrolled by the writers in the central part of the state contained less than six mergansers per mile. These

observations lead to the conclusion that when ice formation drives mergansers from the lakes they tend to settle on the first suitable open water to present itself, with the result that streams in the middle of the state are affected much less than those near the lakes. While the majority of the trout water is likely to be relatively free from mergansers, there may exist local areas of dangerous concentration.

On most of the streams which remained at least partly open, American Goldeneyes were present in large numbers. Previous work done by the Institute (unpublished) has shown that the diet of the Goldeneye under such circumstances is made up almost wholly of roots of submerged vegetation, and of crayfish, insect larvae, and snails.

Table 1
Analysis of Merganser Stomach Contents

| | Brown Trout | Rainbow Trout | Brook Trout | Trout sp. | Suckers | Perch | Black Bass | Other Fish | Crayfish | Frogs | Insects | Debris |
|--|-------------|---------------|-------------|-----------|---------|-------|------------|------------|----------|-------|---------|--------|
| Number of birds containing | 20 | 21 | 8 | 11 | 6 | 9 | 1 | 27 | 18 | 4 | 5 | 21 |
| Average length in inches (to nearest 0.1 inch) | 8.6 | 6.0 | 6.2 | 4.1 | 4.4 | 2.8 | 4.0 | 2.5 | | | | |
| Average number per stomach | 0.48 | 0.64 | 0.09 | 0.15 | 0.19 | 0.39 | 0.05 | 0.61 | 0.20 | 0.04 | 0.06 | |
| Per cent, of total contents based on measured volume | 47.4 | 22.5 | 4.9 | 2.8 | 8.8 | 3.3 | 1.0 | 5.1 | 2.1 | 0.8 | 0.4 | 0.9 |

Explanation of table: All figures based on analysis of 100 stomachs, 98 of American merganser, 2 of Red-breasted Merganser. Under the heading "suckers" are included 18 Common Suckers (*C. commersonii*), and 1 Red-horse, (*Moxostoma* sp.). Under the heading "other fish" are included lamprey ammocoete, Rock Bass, Common Sunfish, Log Perch, Black-sided Darter, Menona Killifish, Black-nosed Dace, Lake Emerald Shiner, various unidentifiable minnows of the genus *Notropis*, Mud Minnow and Muddler. The insects include 2 phryganeid caddis larvae, a giant water bug (*Lethocerus*), and 3 dytiscid beetles. The frogs are apparently all the green frog, *Rana clamitans*.

Notes: We wish to express our thanks to our colleague in the Institute, Mr. Gerald P. Cooper, for determination of the non-game fish.

Of the 100 stomachs examined, 56 contained trout; 33 had eaten trout exclusively.

The average number of trout for each of the 56 stomachs was approximately 2.4.

Table 2

Distribution of Merganser Collections, Trout and Non-trout Waters

| Date (1936) | Locality | Trout or Non-trout | Number of Mergansers | No. containing trout |
|----------------|--|--------------------|----------------------|----------------------|
| Feb.17-Mar.8. | Platte River, Benzie County | Trout | 36 | 28 |
| Feb.21-Mar.1. | Boardman River, Grand Traverse County, | Trout | 13 | 4 |
| Feb.26-29. | South Branch Au Sable River, Crawford County, | Trout | 9 | 8 |
| Feb.24-Mar.17. | West Arm of Grand Traverse Bay, Grand Traverse County, | Non-trout | 7 | 0 |
| Feb. 23-25. | Muskegon River, Newaygo County, | Trout | 5 | 1 |
| Feb. 18-25. | Baldwin Creek, Lake County, | Trout | 5 | 2 |
| Feb.17-Mar.19. | Big Betsie River, Benzie County, | Trout | 4 | 1 |
| Feb. 22. | East Branch Big Creek, Oscoda County, | Trout | 4 | 3 |
| Feb. 23. | Main Stream of Au Sable River, Oscoda County, | Trout | 3 | 2 |
| Feb.19-Mar.21. | North Branch Au Sable River, Crawford County, | Trout | 3 | 1 |
| Feb. 23. | Middle Branch River, Osceola County, | Trout | 3 | 1 |
| Feb.27-29. | Hatcheries, Van Buren County, | Trout | 2 | 2 |
| Feb. 21. | White River, Newaygo County, | Trout | 2 | 0 |
| Feb.28-Mar.1. | Houghton Lake, Roscommon County, | Non-trout | 2 | 0 |
| Feb. 29. | Hatchery, Marquette County | Trout | 1 | 1 |
| Feb. 29. | Pine River, Wexford County, | Trout | 1 | 0 |

From the above table it may be seen that nine of the ducks were taken on non-trout water. Considering only those taken on trout water the percentage containing trout of all species becomes 61.04, almost two-thirds of the total. In certain instances the figure will be seen to be much higher.

It may be of interest to mention that during the peak of the merganser concentration the authors carried on extensive seining operations in the North Branch of the Au Sable, and secured almost no trout, although considerable numbers of Black-nosed Dace, Horned Dace and Muddlers were taken.

It should be borne in mind that the period of heavy merganser concentration on trout streams seldom exceeds thirty days. However, although the figures listed above would have more significance were more known about the merganser's rate of digestion, they do strongly hint that occasional local centers of merganser abundance may reach a strength definitely harmful to the trout population.